

UCRL- 94779
PREPRINT

MODEL ATMOSPHERES FOR X-RAY BURSTING
NEUTRON STARS

Richard A. London, Ronald E. Taam and
W. Michael Howard
University of California
Lawrence Livermore National Laboratory
Livermore, CA 94550

CIRCULATION COPY
SUBJECT TO RECALL
IN TWO WEEKS

This paper was prepared for submittal to the
International Astronomical Union
Symposium No. 125
Nanjing, China
May 25-30, 1986

June 1986

Lawrence
Livermore
National
Laboratory

This is a preprint of a paper intended for publication in a journal or proceedings. Since changes may be made before publication, this preprint is made available with the understanding that it will not be cited or reproduced without the permission of the author.

DISCLAIMER

This document was prepared as an account of work sponsored by an agency of the United States Government. Neither the United States Government nor the University of California nor any of their employees, makes any warranty, express or implied, or assumes any legal liability or responsibility for the accuracy, completeness, or usefulness of any information, apparatus, product, or process disclosed, or represents that its use would not infringe privately owned rights. Reference herein to any specific commercial products, process, or service by trade name, trademark, manufacturer, or otherwise, does not necessarily constitute or imply its endorsement recommendation, or favoring of the United States Government or the University of California. The views and opinions of authors expressed herein do not necessarily state or reflect those of the United States Government or the University of California, and shall not be used for advertising or product endorsement purposes.

MODEL ATMOSPHERES FOR X-RAY BURSTING NEUTRON STARS*

Richard A. London, Ronald E. Taam and W. Michael Howard
Physics Department, LLNL
University of California
P.O. Box 808
Livermore, CA 94550 U.S.A.

Self consistent neutron star atmospheric models have been constructed which include the effects of Comptonization, free-free and bound-free absorption. It has been demonstrated that for parameters relevant to x-ray bursting neutron stars the atmosphere does not radiate like a blackbody during any phase of an x-ray burst. In particular, during the initial rise and final decline of the burst the temperature structure of the atmosphere is affected by backwarming associated with the high opacity due to free-free processes at low frequencies to an extent that the radiation spectrum is shifted to higher energies than a blackbody of the same effective temperature. On the other hand, near the peak of the burst, the opacity is more gray-like as the electron scattering opacity dominates; however, in this case thermalization of the radiation field occurs at such large optical depths ($\tau \sim 5$) that the spectral temperature is higher than the effective temperature. This result is found despite the importance of Comptonization in the thermalization process. Thus, the super Eddington fluxes implied by the spectral data alone are misleading and result from the improper use of the spectral temperature for the effective temperature. For neutron stars characterized by a soft equation of state and radiating near the Eddington effective temperature, fluxes obtained in this way could be overestimated by a factor of about 5.

Because the spectral hardening factor varies throughout an x-ray burst, deconvolution of the spectrum is required before definitive statements can be made concerning the variation of the size of the emitting region. We may say that because of the spectral hardening effect, radius determinations, based on the spectral temperature are only lower limits.

We have also found that the shape of the continuum can be affected by the presence of Fe in the atmosphere; however, the features from bound-free transitions in partially ionized Fe are not particularly strong in the non-LTE approximation for solar abundances.

*Work performed under the auspices of the U.S. DOE by the LLNL under contract number W-7405-ENG-48.